



**Phylogeny of *Himantormia* - an Antarctic genus in the Parmeliaceae (lichenized ascomycetes).**

Thell, A.; Søchting, Ulrik; Kärnefelt, I.; Elix, J. E.; Sancho, L. G.

*Published in:*  
Bibliotheca Lichenologica

*Publication date:*  
2007

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Thell, A., Søchting, U., Kärnefelt, I., Elix, J. E., & Sancho, L. G. (2007). Phylogeny of *Himantormia* - an Antarctic genus in the Parmeliaceae (lichenized ascomycetes). *Bibliotheca Lichenologica*, 95, 531-541.

- POELT, J. (1970): Das Konzept der Artenpaare bei den Flechten. – *Vorträge aus dem Gesamtgebiet der Botanik hrsg. von der deutschen botanischen Gesellschaft, Berlin* **4**: 187–198.
- POELT, J. (1972): Die taxonomische Behandlung von Artenpaaren bei den Flechten. – *Botaniska notiser* **125**: 77–81.
- TEHLER, A. (1983): The genera *Dirina* and *Roccellina* (Roccellaceae). – *Opera Botanica* **70**: 1–86.
- TEHLER, A. (1990): A new approach to the phylogeny of Euascomycetes with a cladistic outline of Arthoniales focussing on Roccellaceae. – *Canadian Journal of Botany* **68**: 2458–2492.
- TEHLER, A. (2002): *Roccella*. – In: NASH III, T.H., RYAN, B., GRIES, C. & BUNGARTZ, F. (eds.), *Lichen Flora of the Greater Sonoran Desert Region. Volume I (the pyrenolichens and most of the squamulose and macrolichens). Lichens unlimited*, Tempe, Arizona, USA.
- TEHLER, A. (2006): *Roccella gracilis* the correct name for *R. peruensis* and *R. humboldtiana* (lichenized fungi, Roccellaceae). – *Taxon* **55**(3): 785–787.
- TEHLER, A., DAHLKILD, Å., ELDENÄS, P. & FEIGE, G.B. (2004): The phylogeny and taxonomy of Macaronesian, European and Mediterranean *Roccella* (Roccellaceae, Arthoniales). – *Symbolae Botanicae Upsalienses* **34**(1): 405–428.
- TEHLER, A., LOHTANDER, K., MYLLYS, L. & SUNDIN, R. (1997): On the identity of the genera *Hubbsia* and *Reinkella* (Roccellaceae). – *Symbolae Botanicae Upsalienses* **32**(1): 255–265.
- WEBER, W.A. (1986): The lichen flora of the Galapagos Islands, Ecuador. – *Mycotaxon* **27**: 451–497.
- WEBER, W.A. & BECK, H. (1985): Effects on cryptogamic vegetation (lichens, mosses and liverworts). – In: ROBINSON, G. & DEL PINO, E.M. (eds.), *El Niño en las Islas Galapagos: el evento de 1982-1983*. Fundación Charles Darwin para las Islas Galápagos, Quito.
- WERNER, B. (2000): Untersuchungen zur Artbildung bei strauchigen Roccellaceen (Arthoniales) auf den tropisch-pazifischen Galápagos-Inseln (Dissertation). – Mathematisch-Naturwissenschaftliche Fakultät der Universität zu Köln, Köln.

*Lichenological Contributions in Honour of David Galloway.*

I. Kärnefelt & A. Thell (eds.): *Bibliotheca Lichenologica* **95**: 531–541.

J. Cramer in der Gebrüder Borntraeger Verlagsbuchhandlung, Berlin · Stuttgart, 2007.

## Phylogeny of *Himantormia* – an Antarctic genus in the Parmeliaceae (lichenized ascomycetes)

Arne THELL<sup>1</sup>, Ulrik SØCHTING<sup>2</sup>, Ingvar KÄRNEFELT<sup>1</sup>, John A. ELIX<sup>3</sup> & Leopoldo G. SANCHEZ<sup>4</sup>

<sup>1</sup> The Biological Museums, Lund University, Östra Vallgatan 18, SE-223 61 Lund, Sweden; e-mails: Arne.Thell@botmus.lu.se, Ingvar.Karnefelt@botmus.lu.se

<sup>2</sup> Department of Mycology, Botanical Institute, University of Copenhagen, Ø. Farimagsgade 2D, DK-1353 Copenhagen, Denmark; e-mail: ulriks@bi.ku.dk

<sup>3</sup> Department of Chemistry, Faculty of Science, Australian National University, Canberra, A. C. T. 0200, Australia; e-mail: John.Elix@anu.edu.au

<sup>4</sup> Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense de Madrid, E-28040 Madrid, Spain; e-mail: sancholg@eucmax.sim.ucm.es

**Abstract:** The phylogeny of the Patagonian-Antarctic genus *Himantormia* is investigated with nuclear ITS rDNA sequences in PAUP. The monotypic genus *Nimisia* is synonymized with *Himantormia*, and the new combination *H. deusta* is proposed. The secondary chemistry was investigated using HPLC; protocetraric, stictic, succinprotocetraric and confumarprotocetraric acids were detected in *H. deusta* for the first time, and alectorialic, hypoalectorialic and decarboxyhypoalectorialic acids are reported as new for *H. lugubris*.

## Introduction

The monotypic genus *Nimisia* Kärnefelt & A. Thell was described from Tierra del Fuego based on material collected by P. L. Nimis and J. Poelt (KÄRNEFELT & THELL 1993) and the new species, *N. fuegiae* Kärnefelt & A. Thell, was distributed in Vězda's exsiccata a few years later. However, during a project to incorporate the lichen collections of Michigan State University in a data base, older samples of *Nimisia fuegiae* collected by H. Imshaug were found. The combination *Cetraria deusta* (Hook. f. & Taylor) Imshaug remained unpublished, because Imshaug was convinced that it did not belong in *Cetraria s. str.* (FRYDAY 2005). The basionym *Parmelia enteromorpha* Ach. var. *deusta* Hook. f. & Taylor, had earlier been recombined and raised to species level as *Hypogymnia deusta* (Hook f. & Taylor) C. W. Dodge (DODGE 1965). FRYDAY (2005) corrected the name to *Nimisia deusta* (Hook. f.) Fryday and presented new data on its morphology, secondary chemistry and phytogeography of this species. In particular, he found that the outer surface was not entirely black but had

pale brown patches close to the base, that fumarprotocetraric acid was present in the upper medulla, and, that the distribution range included the Cape Horn and Falkland Islands (FRYDAY 2005: 317, map). *Esslingeriana idahoensis* (Essl.) Hale & M. J. Lai and *Himantormia lugubris* (Hue) I. M. Lamb are two other species in the Parmeliaceae which have been compared to *Nimisia* because of similarities in their cortical anatomy, i.e. paraplechtenchyma composed of large dark pigmented cells with comparatively large lumina. *Himantormia lugubris* showed additional similarities with *Nimisia*, such as the pachydermatous medulla and its Antarctic distribution (KÄRNEFELT & THELL 1993). The similar distribution and the absence of lichenan in the cell-walls made *Himantormia* the strongest candidate among possibly related genera (COMMON 1991; FRIDAY 2005). Three additional species with alternative northern hemisphere distributions, *Arctocetraria nigricascens* (Nyl.) Kärnefelt & A. Thell, *Cornicularia normoerica* (Gunnerus) Du Rietz, and *Kaernefeltia merrillii* (Du Rietz) A. Thell & Goward, were discussed as possibly being related to *Nimisia*, because of similarities in overall morphology (KÄRNEFELT & THELL 1993),.

Extensive phylogenetic analyses of the Parmeliaceae, based on DNA-sequences have yet to include *Nimisia*, because of the absence of fresh material, and the fact that the presumably related species mentioned above did not show close affinities with one another (THELL et al. 2004). Fortunately, fresh material of *Nimisia deusta* needed for a DNA-investigation was collected by U. Söchting and L. Sancho during an excursion to the Navarino Island, Southern Chile in 2005.

## Material and methods

### Selected material

The 30 samples, representing 27 species, were selected to represent a range of Parmeliaceae focussing on species presumed to be related to *Nimisia deusta*. The majority of the Parmeliaceae has been divided into four large monophyletic groups, the cetrarioid, parmotremond, usneoid and xanthoparmelioid lichens (OHMURA 2002; THELL et al. 2002; 2004; BLANCO et al. 2004, 2005). These groups are represented in the analysis. Five species possibly related to *Nimisia* were also included (KÄRNEFELT & THELL 1993; FRYDAY 2005). Finally, a blast(n) search was performed, matching the new *Nimisia* sequence in the GenBank, to detect species with high percentage of identities. Five of the sequences are new, whereas 25 were downloaded from the GenBank at [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) (Table 1). A blast(n) search was performed matching the new *N. deusta* sequence with sequences present in the GenBank (ALTSCHUL et al. 1997).

Two cetrarioid species, *Nephromopsis pseudocomplicata* and *N. stracheyi* were selected as the outgroup. The monophyletic group of cetrarioid lichens, investigated in several DNA-analyses, is presumed not closely related to *Nimisia*.

## DNA analysis

**Extraction:** Employing MagAttract 96 DNA Plant Extraction Kit from Qiagen (08/2003), samples were rigorously shaken in a vortex machine in 300 ml extraction buffer incorporating one steel bead into each microtube; otherwise, the enclosed protocol for manual DNA purification was followed.

**Table 1.** Selected specimens with extraction numbers, sample-IDs and GenBank accession numbers.

Species	Extr	Sample-ID	GenB acc
<i>Alectoria ochroleuca</i>	976	AUSTRIA. Ötztal, 2001-08-11, Feuerer & Thell (HBG)	AF451735*
<i>Arctocetraria andrejevii</i>	1364	GREENLAND. Hansen exs. 836 (LD 1001631)	DQ004575*
<i>Arctocetraria nigricascens</i>	793	CANADA. Melville Isl., Westberg 1614 (LD)	AF254628*
<i>Bryocaulon divergens</i>	1914	SWEDEN. Härjedalen, Arup L02258 (LD)	DQ395287
<i>Bryoria fuscescens</i>	920	ITALY. Trentino-Alto Ad., Feuerer & Thell 62282 (HBG)	AF451736*
<i>Bryoria sp.</i>	1932	GREENLAND. Narsarsuaq, Thell & Feuerer (LD 1039864)	DQ395288
<i>Coelopogon abraxas</i>	1209	CHILE. 2001-11-11, Feuerer (HBG)	AY251414*
<i>Cornicularia normoerica</i>	930	ITALY. Trentino-Alto Ad., 2001, Feuerer & Thell (HBG)	AY251416*
<i>Dactylina arctica</i>	160	CANADA. Alberta, Miao (TDI 300)	AF115760*
<i>Esslingeriana idahoensis</i>	146	CANADA. B. C., Goward 961348 (UBC)	AF227513*
<i>Himantormia deusta</i>	1910	CHILE. Region XII Magallanes, Söchting 10257 (C)	DQ395290
<i>Himantormia lugubris</i>	1231	ANTARCTICA. South Shetland Islands, Söchting 7609 (C)	AF251421*
<i>Himantormia lugubris</i>	----	ANTARCTICA. Kim 05013	DQ219309*
<i>Hypogymnia physodes</i>	16	SWEDEN. Skåne, Thell 9605 (LD 1029842)	AF141368*
<i>Kaernefeltia californica</i>	1703	U. S. A. Oregon, McCune 27703 (LD 1045103)	DQ004571*
<i>Kaernefeltia merrillii</i>	190	CANADA. British Columbia, Thell 9698 (LD 1089087)	AF072230*
<i>Kaernefeltia merrillii</i>	1916	SPAIN. Madrid, El Berrueco, Thell et al. (LD 1044393)	DQ395291

<i>Kaernefeltia merrillii</i>	1918	SPAIN. Madrid, El Berruoco, Thell et al. (LD 1038537)	DQ395292
<i>Lethariella intricata</i>	----	SPAIN. Chanary Islands, Kroken & Taylor (2000)	AF297742*
<i>Melanelia stygia</i>	922	ITALY. Trentino-Alto Ad., Feuerer & Thell 64247 (HBG)	AF451775*
<i>Menegazzia terebrata</i>	916	GERMANY. Bayern, 2001, Feuerer & Thell (HBG)	AY251430*
<i>Nephromopsis pseudocomplicata</i>	907	CHINA. Sichuan, Obermayer 8276a (GZU)	AF404131*
<i>Nephromopsis stracheyi</i>	606	BHUTAN. Thimpu, Søchting 8095 (LD)	AF451785*
<i>Nodobryoria abbreviata</i>	45	CANADA. B. C., Thell & Veer 9645b (LD 1192360)	AF116177*
<i>Parmelia saxatilis</i>	518	CHILE. Region XII Magallanes, Feuerer 29542 (HBG)	AF410672*
<i>Parmotrema chinense</i>	918	GERMANY. Bavaria, 2001, Feuerer & Thell (HBG)	AF451749*
<i>Pseudephebe minuscula</i>	931	ITALY. Trentino-Alto Ad., 2001, Feuerer & Thell (HBG)	AY251446*
<i>Pseudephebe pubescens</i>	884	CHILE. Valdivia, Feuerer 29546 (HBG)	AF451738*
<i>Pseudevernia furfuracea</i>	781	FINLAND. Regio Ab., Feuerer & Thell (LD 1057234)	AF451768*
<i>Usnea florida</i>	840	SWEDEN. Skåne, Thell 0011 (TUR)	AF451739*
<i>Xanthoparmelia pulla</i>	935	AUSTRIA. Tyrol, Feuerer & Thell 64210 (HBG)	AF451747*

**Amplification:** 20 µl PCR-reactions were prepared to amplify the nuclear ITS1-5.8S-ITS2 ribosomal DNA region. The primers ITS1F (GARDES & BRUNS 1993) and ITS4 (WHITE 1990) were used. Each reaction included 5 µl water, 2 µl buffer, 8 µl nucleotides, 1 µl of the extracted DNA, and 0.1 µl of the DNA polymerase. The PCR programme described by EKMAN (2001) was employed: after a 2 min. hold at 94°C, six cycles followed with denaturation at 94° for 60 sec., annealing at 62°-56°C for 60 sec. (decreasing 1° per cycle), and extension at 72°C for 105 sec., followed by 34 cycles with denaturation at 94° for 30 sec., annealing at 56°C for 30 sec. and extension at 72°C for 105 sec.; finally, a 10 min. hold at 72°C was performed before the PCR-products were cooled to 4°C.

**Purification and sequencing:** The PCR-products were cleaned with PCR clean-up NucleoFast 96 PCR Purification Kit from Macherey-Nagel as per the enclosed user manual (2002/3/Rev. 01. p.12). 50 µl TE buffer was applied to each sample prior to the standard procedure for purification of PCR-products under vacuum, which was followed, except for the second, optional, washing step, with purified water. The purified samples were

collected in 50 µl water and the amount of DNA was measured in an Eppendorf BioPhotometer. The concentrations differed between 18-54 ng/µl. The required amount of DNA for sequencing, 1 ng/base pair, was dried for 1 hr at 65°C. Finally, the DNA-fragments were sent to Macrogen to be sequenced, again using the primers ITS1F and ITS4.

**Phylogeny:** The phylogenetic analyses of the manually aligned ITS sequences were done with PAUP version 4.0b (SWOFFORD 1998). Trees were calculated using the general heuristic search option. Gaps were treated as missing characters. Bootstrap analyses with 1000 replicates were performed using the same settings. Support values of 50 or above are marked in the consensus tree (Fig. 1).

### Anatomy

Most data concerning the morphology and anatomy was obtained from earlier publications on *Himantormia* (LAMB 1964; KÄRNEFELT & THELL 1993; FRYDAY 2005), but some new anatomical sections were made to complement this data. Thalli and apothecia were sectioned (15 µm thick) using a Kryomate, Leitz freezing microtome and conserved in lactophenol cotton-blue. The sections were studied with a Zeiss Axioscope light microscope.

### Secondary chemistry

Secondary compounds were detected by means of high performance liquid chromatography (ELIX et al. 2003).

### Results

#### Blast(n) search

Performing a blast(n) search (ALTSCHUL et al. 1997) with the *Nimisias deusta* sequence in the GenBank revealed the highest percentage of matches together with two *Himantormia lugubris* sequences, DQ219309 and AF251421, showing differences at c. 20 and 21 positions (98%) respectively.


#### Phylogeny analysis

The aligned matrix was composed of 523 characters, of which 162 were parsimony informative. The heuristic search resulted in a single most parsimonious tree, 757 steps long, CI=0.4848, RI=0.4855, RC=0.2354 (Fig. 1). The bootstrap analysis, using the same preferences as in the heuristic search, showed maximum support for a common clade for *Nimisias deusta* and *Himantormia lugubris*.

## *Himantormia* I. M. Lamb

*Brit. Antarctic Surv. Sci. Rep.* **38**: 17-18 (1964). – **Typus generis:** *Himantormia lugubris* (Hue) I. M. Lamb.

**Syn.:** *Nimis* Kärnefelt & A. Thell, *Lichenologist* **25**: 370–371 (1993).



**Fig. 1.** The phylogenetic position of *Himantormia* in the Parmeliaceae. The single most parsimonious tree length = 757, CI = 0.4848, RI = 0.4855, RC = 0.2354. Paup 4.0b. Bootstrap support values of 50 or above from 1000 replicates are indicated at the branches.

The hitherto monotypic genus *Himantormia* is now proposed to comprise two species, both of which have a Patagonian-Antarctic distribution. Thus a new combination is proposed for the sole species previously accommodated in the genus *Nimis*. Both species have previously been described in detail

(LAMB 1964, KÄRNEFELT & THELL 1993), and data presented below mainly concerns new discoveries in phytogeography and secondary chemistry.

## *Himantormia deusta* (Hook. f.) A. Thell & Søchting, *comb. nov.* (Fig. 2)

**Bas.:** *Parmelia enteromorpha* var. *deusta* Hook. f., *Flora Antarctica* (Bot. Antarct. Voyage Erebus & Terror 1): 532 (1847). – *Hypogymnia deusta* (Hook. f.) C. W. Dodge, *Trans. Amer. Microscop. Soc.* **84**: 506 (1965). – *Cetraria deusta* (Hook. f.) Imshaug, in sched. – **Typus:** CHILE. Tierra del Fuego, near Cape Horn, c. 56°30'S, 60°W, Mt Foster, ix 1842, J. D. Hooker, *Voyage Erebus & Terror* (E – lectotype, Friday, *Lichenologist* **34**: 315, 2005).

**Syn.:** *Nimis fuegiae* Kärnefelt & A. Thell, *Lichenologist* **25**: 370–371 (1993). – **Typus:** ARGENTINA. Isla Grande de Tierra del Fuego, Dept. Ushuaia, Sierra Alvear, just in front of Hosteria Tierra Major, on steeply inclined faces of granite rocks in the subalpine belts, 1987, P. L. Nimis & M. Tretiach (LD – holotype; TSB, DUKE – isotypes; Vezda: Lichenes Rariores Exsiccati, no 144; TSB 11131; leg. P. L. Nimis, M. Codogno & M. Tretiach).

The spermatia of *H. deusta* are almost rod-shaped but slightly thicker in the middle, hence the reason for classifying them as oblong citriform by KÄRNEFELT & THELL (1993); however, their size and shape are in fact very similar to the bacillariform spermatia of *H. lugubris* (Table 2).

Although this species was originally reported to have no chemical compounds (KÄRNEFELT & THELL 1993), FRYDAY (2005) detected fumarprotocetraric acid whereas fumarprotocetraric (major), protocetraric (minor), stictic (trace), succinprotocetraric (trace) and confumarprotocetraric (trace) acids have been detected by HPLC in the present study, none of which occur in *H. lugubris*.

## *Himantormia lugubris* (Hue) I. M. Lamb

(Fig. 3)

*Brit. Antarctic Survey Sci. Rept* **38**: 18–23 (1964). – **Bas.:** *Ramalina lugubris* Hue, *Exp. Antarct. Franc.*: 34–36 (1915). – **Typus:** Antarctica. Ile Petermann, peu commun sur les rochers du sommet de l'île, altitude 120 mètres, XXe excursion, no 286, le 14 mars 1909, Hue (PC – holotype).

The excellent description of *H. lugubris* made by HUE (1915) and LAMB (1964) is completed by the detection of additional secondary compounds using modern HPLC-equipment. However, the anatomy of both *Himantormia* species was reinvestigated, the distinguishing characters of which are presented in Table 2.

LAMB (1964) reported small quantities of atranorin and barbatolic acid as an unidentified b-orcinol derivative and HUNECK & FOLLMANN (1966) detected usnic acid in addition to these two secondary compounds. Alectorialic (major), hypoalectorialic (minor), barbatolic (trace) and decarboxyhypoalectorialic (trace) acids have been detected by HPLC in the present study.



Fig. 2. *Himantormia deusta*, x 5, US 10257 (C).



Fig. 3. *Himantormia lugubris*, x 5, US 7609 (C).

## Discussion

In a recent survey, *Himantormia lugubris* was found to belong to a section of the Parmeliaceae composed mainly of isolated genera and small groups of genera without any evident relationships with one another (THELL et al. 2002; 2004). Blast(n) searches for matching sequences with four additional two-species genera in the analysis show similar amount of matches between the species compared with *Himantormia*, where 96% matches were

detected. In *Arctocetraria* Kärnefelt & A. Thell, the blast(n) search resulted in approximately the same rate of matches between the species, *A. andrejevii* and *A. nigricascens*, although the bootstrap support of 82 % was weaker (Fig. 1). Of the genera *Kaernefeltia* A. Thell & Goward and *Pseudephebe* Choisy, the latter has the highest amount of matches, 98%, when performing a blast(n) search and, in similarity with *Himantormia*, a 100% support in the bootstrap analysis. *Kaernefeltia* has 97-98% of matches. Notably, the variation between western American and Spanish populations of the disjunct species *K. merrilli* were larger than that between *K. merrillii* and the second species of the genus, *K. californica*. The genus has a moderate bootstrap support of 73% in the phylogeny analysis (Fig 1, cf. THELL et al. 2005: Figs. 1-3).

Table 2. A comparison of diverging characters of *Himantormia deusta* and *H. lugubris* (Figs. 2-3).

Character	<i>Himantormia deusta</i>	<i>Himantormia lugubris</i>
Lobes, shape and height	strap-shaped, slightly canaliculate, to 1 cm	awl-shaped to strap-shaped, to 4 cm
Apothecia, position	laminal	laminal, rarely terminal
Cortex, cell size	10–15 µm	4–7 µm
Medulla, structure	dense	lax, with dense portions
Medulla, hyphae diam.	7–9 µm	4–7 µm
Medulla, hyphae lumina diam.	0.1–0.5 µm	1–2 µm
Algae, position	below pycnida and apothecia	below cortex in light pigmented patches of areolae or verrucae
Spermatia, shape	almost bacilliiform, slightly thicker in the middle	bacilliiform
Spermatia, size	8–10 x 1–1.5 µm	9–10 x 0.8–1 µm
Secondary compounds	fumarprotocetraric acid (major), protocetraric acid (minor), stictic acid (trace), succinprotocetraric acid (trace) and confumarprotocetraric acid (trace)	alectorialic acid (major), hypoalectorialic acid (minor), barbatolic acid (trace) and decarboxy-hypoalectorialic acid (trace) by HPLC. Atranorin (reported by LAMB (1964), minor).

## Acknowledgements

Prof. Mark Seaward is thanked for correcting the English and useful comments on a draft of this paper. The DNA-investigations were performed by the first author in Copenhagen, in January 2006, within the SYNTHESYS project DK-TAF U1412. Support from the Carlsberg Foundation to the second author is acknowledged.

## References

- ALTSCHUL, S.F., MADDEN, T.L., SCHÄFFER, A.A., ZHANG, J., ZHANG, Z., MILLER, W. & LIPMAN, D.J. (1997): Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. – *Nucleic Acids Research* **25**: 3389–3402.
- BLANCO, O., CRESPO, A., DIVAKAR, P.K., ELIX, J.A. & LUMBSCH, T.H. (2005): Molecular phylogeny of parmotreoid lichens (Ascomycota, Parmeliaceae). – *Mycologia* **97**: 150–159.
- BLANCO, O., CRESPO, A., ELIX, J.A., HAWKSWORTH, D.L. & LUMBSCH, T.H. (2004): A molecular phylogeny and a new classification of parmelioid lichens containing *Xanthoparmelia*-type lichenan (Ascomycota: Lecanorales). – *Taxon* **53**: 959–975.
- COMMON, R.S. (1991): The distribution and taxonomic significance of lichenan and isolichenan in the Parmeliaceae (lichenized Ascomycotina), as determined by iodine reactions. I. Introduction and methods. II. The genus *Alectoria* and associated taxa. – *Mycotaxon* **41**: 67–112.
- DODGE, C.W. (1965): Lichenological notes on the flora of the Antarctic Continent and the Subantarctic Islands – V. Tierra del Fuego and the Falkland Islands. – *Transcripts of the American Microscopical Society* **84**: 502–507.
- EKMAN, S. (2001): Molecular phylogeny of the Bacidiaceae (Lecanorales, lichenized Ascomycota). – *Mycological Research* **105**: 783–797.
- ELIX, J.A., TØNSBERG, T. & WARDLAW, J.H. (2005): The structure of friesiic acid, a novel lichen substance from *Hypocenomyce friesii*. – In: DÖBBELER, P. & RAMBOLD, G. (eds), Contributions to Lichenology. Festschrift in honour of Hannes Hertel. – *Bibliotheca Lichenologica* **88**: 103–109.
- FRYDAY, A.M. (2005): *Nimisia deusta*, the correct name for *N. fuegiae*, with additional notes on morphology, chemical composition, and distribution. – *Lichenologist* **37**: 313–319.
- FRYDAY, A.M. & PRATHER, L.A. (2001): The lichen collection of Henry Imshaug at Michigan State University Herbarium (MSC). – *Bryologist* **104**: 464–467.
- HUE, A.M. (1915): Lichens. – In: *Expédition antarctique française (1903–1905) commandée par le Dr. Jean Charcot*. Paris, 202 pp.
- GARDES, M. & BRUNS, T.D. (1993): ITS primers with enhanced specificity for basidiomycetes. Application to the identification of mycorrhizae and rusts. – *Molecular Ecology* **2**: 113–188.
- KÄRNEFELT, E.I. & THELL, A. (1993): *Nimisia fuegiae* (Parmeliaceae), a new lichen genus and species from Argentina. – *Lichenologist* **25**: 369–377.
- KROKEN, S. & TAYLOR, J.W. (2000): Phylogenetic species, reproductive mode, and specificity of the green alga *Trebouxia* forming lichens with the fungal genus *Letharia*. – *Bryologist* **103**: 645–660.
- LAMB, I.M. (1964): Antarctic lichens I. The genera *Usnea*, *Ramalina*, *Himantormia*, *Alectoria*, *Cornicularia*. – *British Antarctic Survey Scientific Reports* **38**: 1–34.
- LEGAS, M.-E., MATEOS, J.L. & VICENTE, C. (1999): Effect of plasmidic DNA-intercalating acridine orange on the production of barbatolic acid by the lichen *Himantormia lugubris* (Hue) Lamb. – *Journal of the Hattori Botanical Laboratory* **86**: 221–233.
- OHMURA, Y. (2002): Phylogenetic evaluation of infrageneric groups of the genus *Usnea* based on ITS regions in rDNA. – *Journal of the Hattori Botanical Laboratory* **92**: 231–243.
- SWOFFORD, D.L. (1998): PAUP: Phylogenetic analysis using parsimony, version 4.0b. Sinauer Associates, Sunderland.
- THELL, A., FEUERER, T., KÄRNEFELT, I., MYLLYS, L. & STENROOS, S. (2004): Monophyletic groups within the Parmeliaceae identified by ITS rDNA, b-tubulin and GAPDH sequences. – *Mycological Progress* **3**: 297–314.
- THELL, A., RANDLANE, T., SAAG, A. & KÄRNEFELT, I. (2005): A new circumscription of the lichen genus *Nephromopsis* (Parmeliaceae, lichenized ascomycetes). – *Mycological Progress* **4**: 303–316.
- THELL, A., STENROOS, S., FEUERER, T., KÄRNEFELT, I., MYLLYS, L. & HYVÖNEN, J. (2002): Phylogeny of cetrarioid lichens (Parmeliaceae) inferred from ITS and b-tubulin sequences, morphology, anatomy and secondary chemistry. – *Mycological Progress* **1**: 335–354.